



Gynecomastia Instructions

BEFORE YOUR SURGERY:

Suggestions:

- Arrange for 5-7 days off work. Recovery is variable, depending on your individual response to surgery, but some time to heal is mandatory.
- Try to complete all non-essential personal duties and responsibilities before your surgery. Your recovery will be far less stressful if you can get all of life's little tasks out of the way beforehand.
- Have someone available for transportation to and from the hospital.
- Obtain a large package of 4"x4" gauze for use after surgery

Mandatory:

- Stop intake of all Aspirin and Aspirin-related products (anti-inflammatory like Motrin, Advil, Naprosyn, etc.) two weeks before surgery, as these products will increase bleeding and bruising during surgery
- Stop smoking or nicotine-containing products (Nicorette, nicotine patch) 4-6 weeks before surgery. I would prefer 6 weeks if possible.
- Consume no food, drink or even water after midnight the night before surgery. A full stomach on the morning of surgery will necessitate cancellation of the procedure.
- Discuss any medications, or changes in medications by contacting the office.
- Have someone available to bring you home after surgery (usually later in the afternoon).
- Have someone available to stay with you the night after surgery.
- Be at the hospital before your scheduled time, the hospital will contact you with this exact information.
- Call your surgeon with any questions you may have, as soon as possible, before your surgery.

OTTAWA PLASTIC SURGERY

DR. H. SILVERMAN, MD, FRCSC
DR. KIRSTY BOYD, MD, FRCSC



AFTER YOUR SURGERY:

Immediately After Your Surgery:

- No heavy lifting (more than 10 pounds)
- Sleeping is usually most comfortable on your back, propped up on pillows, but you may sleep on your side if comfortable
- Wear your conforming garment or tensor at all times, to reduce swelling and support the contour of the area treated
- The dressing is to remain dry and in place until I see you for your first post-operative visit. Sponge baths or sitting in the bath are acceptable, but please keep the bandages dry.
- You may eat a regular diet
- No driving until discussed with me
- Take all medications as prescribed:
 - Usually Dilaudid, Tylenol #3 or Percocet, 1 or 2 tablets every 3 to 4 hours **as needed** for pain. This may be supplemented by over the counter anti-inflammatories, such as Motrin, Ibuprofen, Advil or Tylenol.
 - Note that the pain medicine may make you nauseous or constipated. You may use over-the-counter stool softeners, such as Colace.
- Call my office to arrange a follow-up appointment for about 2-5 days after surgery

Things to Watch Out For:

- A rapid increase in pain, swelling and firmness, with severe bruising
- Vomiting after the first 24-48 hours
- High fevers above 38.5 C or 101.5 F
- Pain and swelling in one, or both legs
- **The presence of these above mentioned conditions warrants a call to your surgeon**

Normal Post-Surgical Reactions:

- Pain, bruising, and mild swelling are normal and expected after surgery
- Low grade temperature is often seen early after surgery
- Mild differences in size are common and normal in the early recovery period

OTTAWA PLASTIC SURGERY

DR. H. SILVERMAN, MD, FRCSC
DR. KIRSTY BOYD, MD, FRCSC



Late Post-Operative Period:

- No heavy lifting, weight training, vigorous exercise such as aerobics, jogging or swimming, for 4 weeks after surgery
 - Driving is allowed once you are off narcotic pain medication, **and** can move comfortably enough to react to any driving situation appropriately; check with me if you are unsure
 - You may shower after the first office dressing change, and gently wash directly over the incisions
 - Wear your conforming garment at all times for the first 4 weeks, unless otherwise directed
 - Dry gauze may be placed inside the bra if small amounts of drainage persist
 - Your stitches will be removed approximately two weeks after surgery. Creams and lotions over the scars are not to be used unless otherwise directed
 - Normal activities may be resumed as tolerated by 2-4 weeks after surgery
 - Avoid tanning over healing incisions
-
- **If any of the above information is unclear, or any other questions arise, please do not hesitate to contact our office, during business hours, Monday to Friday – 08:30am-4:30pm**
 - **After hours and weekends: call my office, where you will be directed to call an emergency contact number, which is either Dr. Silverman’s cell phone – 613-355-9007, or Dr. Boyd’s cell phone – 613-297-2974.**
 - **Additionally you can contact us by email directly to askdrsilverman@ottawaplasticsurgery.com or askdrboyd@ottawaplasticsurgery.com, though we cannot guarantee an emergent response via email.**
 - **If you are unable to reach us and have a medical emergency, please go to the nearest emergency room.**